


Transactions NAST PHL, is the official journal of the National Academy of Science and Technology Philippines. It has traditionally published papers presented during the Academy's Annual Scientific Meeting since 1979 to promote science-based policy discussions of and recommendations on timely and relevant national issues as part of its functions as a national science academy. Starting in 2021, this journal has been open to contributions from the global scientific community in all fields of science and technology.



Research Note: Biologically Active Constituents from *Mentha Cordifolia Opiz* Leaves

Irene M. Villaseñor, Jennifer Angelada, Arlyn Canlas,
Deborah Echegoyen

Institute of Chemistry, University of the Philippines
Diliman, Quezon City

Citation

Villaseñor IM et al. 1999. Biologically active constituents from *Mentha Cordifolia Opiz* leaves. Transactions NAST PHL 21: 197-200. doi.org/10.57043/transnastphl.1999.5771

Copyright

© 1999 Villaseñor IM et al

Research Note:

BIOLOGICALLY ACTIVE CONSTITUENTS FROM *MENTHA Cordifolia* OPIZ LEAVES

IRENE M. VILLASEÑOR, JENNIFER ANGELADA,
ARLYN CANLAS, DEBORAH ECHEGOYEN
Institute of Chemistry, University of the Philippines, Diliman, Q.C.

ABSTRACT:

Isolates from yerba buena leaves were tested for their analgesic, antimutagenic, and anthelmintic activities. The acetic acid-induced writhing test showed that isolates β -sitosterol, β -sitosteryl- β -D-glucoside, and cis-8-pentadecenylactone, at a dosage of 100 mg / kg mouse, each decreased the number of squirms induced by acetic acid by 70.0%, 73.0%, and 67.3%, respectively. β -sitosterol and an unsaturated carboxylic acid derivative are antimutagenic because they inhibited the mutagenicity of tetracycline by 65.3% and 68.7%, respectively, at a dosage of 0/01 mg / 20 g mouse, using the *in vivo* Micronucleus Test. *In vitro* tests using *Ascaris suum* showed that β -sitosterol is also anthelmintic as the behavior of worms approximate that of the positive controls, Combantrin and Antiox.


Keywords: *Mentha cordifolia* Opiz., yerba buena, analgesic, anthelmintic, antimutagenic, β -sitosterol, β -sitosteryl- β -D-glucoside

Mentha cordifolia Opiz., commonly known as Philippine mint, marshmint, or yerba buena, is listed as one of the priority plants under the Department of Science and Technology (DOST)-Philippine Council for Health Research and Development (PCHRD)- National Integrated Research Program on Medicinal Plants (NIRPROMP). The unextracted and unpurified leaves are presently being produced in tablet form, including pediatric tablets, and have been proven as an analgesic in clinical trials phases I, II, and III (DOST Technical Report Series No. 12, 1991).

This paper is on the bioassay-directed isolation and structure elucidation of the bioactive constituents from yerba buena leaves. The leaves were extracted by immersing them in methanol. The methanolic extract was then solvent-partitioned into hexane, CHCl_3 , and EtOAc extracts. Subsequent bioassay using the acetic acid-induced writhing test showed that the hexane extract is analgesic. *In-vitro* tests using live *Ascaris suum* as test animals showed that the hexane extract is also anthelmintic. An *in-vivo* Micronucleus Test showed that the CHCl_3 extract is


antimutagenic. The bioactive extract was then purified by sequential and repeated normal phase vacuum liquid chromatography using gradient elution. The structures of the bioactive isolates were elucidated by spectral analysis including gc-msd, ir, ^1H - and ^{13}C -nmr, COSY, HMQC, and HMBC.

The hexane fraction on sequential and repeated vacuum liquid chromatography (vlc) over silica gel afforded the analgesic fractions labeled FB2 and FB10, eluted out at 20% EtOAc/ hexane and 30% EtOH/EtOAc, respectively. Vlc over silica gel of fraction FB2 yielded a greenish crystalline analgesic fraction which on recrystallization using ether/MeOH afforded white needle-like crystals 1 with an R_f 0.76, tlc (silica gel), EtOAc/hexane, 30:70, fuschia spot with vanillin- H_2SO_4 . Spectral analyses gave β -sitosterol.



β -sitosterol

Fraction FB10 on vlc over silica gel 60G afforded 7 fractions with fraction FB10E, eluted from 8% MeOH/ CHCl_3 , exhibiting the highest analgesic activity. Vlc of FB10E yielded white crystals 2 upon recrystallization in ether/MeOH gave with an R_f 0.30, tlc (silica gel), MeOH/ CHCl_3 , 10:90, fuschia spot with vanillin- H_2SO_4 . Spectral analyses gave β -sitosteryl- β -D-glucoside.



β -sitosteryl- β -D-glucoside

Confirmatory bioassay showed that β -sitosterol is analgesic, anthelmintic, and antimutagenic (Fig. 1, 2, 3). Its glucoside is also analgesic.

The analgesic cis-8-pentadecenyllactone is a white crystalline solid which degrades at 135.1°C. The antimutagenic unsaturated carboxylic acid is a yellowish crystalline solid which degrades at 210.8°C.




Figure 1. Analgesic activity using the acetic acid – induced writhing test




Figure 2. Analgesic activity using the hot plate method




Figure 3. Antimutagenicity and mutagenicity using the micronucleus test